



A convenient gel bait insecticide  
for the control of cockroaches in  
residential and industrial situations.

- Easy to use and long lasting
- Cost effective
- Broad Spectrum of Use Locations
- Precise delivery

**Pest Control Solutions**

 **BASF**

The Chemical Company



**Amdro® Gel** is easy to apply with precision in the least possible time, to give long lasting and cost-effective cockroach control.

**Amdro Gel** is great for use in sensitive areas and is labeled for use in commercial situations.

**Amdro Gel** is ideal for use with a non-repellent spray such as **Phantom®** Insecticide for an effective and reliable IPM Program.

## An Easy to Use, Long Lasting Gel

**Amdro® Cockroach Gel** is an easy to use bait insecticide that is effective for long lasting control of cockroaches. The active ingredient hydramethylnon affects the insect's metabolic system, eradicating cockroaches within 24 to 72 hours.

**Amdro Gel** will control all major pest species of cockroaches, including German, Australian and American cockroaches.

**Amdro Gel** lasts up to 3 months through the baiting method thus saves costs, time and product. The reduction of cockroaches can be noticed within a week after application.

## Unique Delayed Action Mortality

**Amdro Gel** is used as a spot or crack-and-crevice treatment, and should only be applied to areas that the cockroaches pass through. Once the cockroaches encounter the palatable gel, they devour it.

Cockroaches who feed on **Amdro Gel** bait do not die immediately. Instead, they return to their harborages carrying the active ingredient with them where other roaches ingest **Amdro Gel** by feeding on their faeces.





## Broad Spectrum of Use Locations

**Amdro Gel** can be used in a large array of both domestic and commercial situations including in and around homes, institutional, warehousing and commercial establishments, including hospitals, laboratories, computer facilities, sewers, buses and other transport vehicles or vessels, pet shops, and zoos.

Gel placements should be at or near harborages or aggregation areas, such as cracks and crevices, corners, or areas of cockroach movement.

Use locations include undersurfaces only of tables, counters, shelving, drawers, and sinks and around pipe collars, above suspended ceilings, inside electrical boxes, adjacent or under trash receptacles and carts, under and behind stoves and refrigerators, under drain plates and around electrical equipment and machines.

## Effective Use for the Greatest Value.



**Amdro Gel** contains 20g/kg Hydramethylnon, in a palatable and stable gel bait. It can be applied in residential and commercial situations and is available in 30g cartridges.

**Amdro Gel** bait will remain palatable to cockroaches as long as it is visibly present. Re-apply **Amdro Gel** when bait is no longer visibly present, according to the remaining level of infestation, which is generally at a lower rate than the initial treatment.

A visual inspection of bait placements is recommended at one week after the initial treatment where high infestations are being treated. Reapplication should be made if the bait is visibly depleted.

For an effective Integrated Pest Management (IPM) program, BASF recommends using a non-repellent spray, such as Phantom<sup>®</sup> Insecticide in conjunction with **Amdro Gel** for a reliable cockroach treatment.

### Amdro Gel in Spot Size Recommendation

Cockroach Species	German Cockroach	Large Cockroach, such as American, Smoky Brown or Oriental
Application Rate (g/m <sup>2</sup> INFESTED SURFACE AREAS)	 0.25 - 0.50g Small Spots	 0.50 - 1.0g Large Spots





### Situations

Residential and commercial situations including in and around homes, institutional, warehousing and commercial establishments, including hospitals, laboratories, computer facilities, sewers, buses and other transport vehicles or vessels, pet shops, and zoos.

### Gel placement

At or near harborages or aggregation areas, such as cracks and crevices, corners, or areas of movement. Use locations include undersurfaces only of tables, counters, shelving, drawers, and sinks and around pipe collars, above suspended ceilings, inside electrical boxes, adjacent or under trash receptacles and carts, under and behind stoves and refrigerators, under drain plates and around electrical equipment and machines.

### Critical comments

Use the lower application rate for low to moderate infestation and the higher rate for more severe infestations

### Restrains

- DO NOT use in areas accessible to children and pets.
- DO NOT use in exposed areas which may be washed.
- DO NOT use near open food and food contact surfaces.
- DO NOT use this product on or around electrical equipment due to the possibility of electrical shock.
- DO NOT apply gel to areas which have been recently sprayed with insecticides, or spray insecticides over gel placements, as this may cause the bait to become repellent.
- DO NOT apply to surfaces where the temperature exceeds 54°C.

## IMPORTANT INFORMATION

THIS IS NOT THE LABEL AND USERS SHOULD REFER TO THE PRODUCT LABEL FOR FULL DETAILS.

BASF Australia Ltd  
Level 12, 28 Freshwater Place, Southbank VIC 3006  
Product Information Hotline  
For further information, contact Freecall 1800 006 393  
[www.pestcontrol.basf.com.au](http://www.pestcontrol.basf.com.au)

Amdro® Gel should be applied by a licensed pest controller only.  
Always read the product label before use.  
This brochure is intended as advice.

Disclaimer: The information submitted in this publication is based on current BASF knowledge and experience. In view of the many factors that may affect its application, this data does not relieve the user from carrying out their own tests. The data does not imply assurance of certain properties or of suitability for a specific purpose. It is the responsibility of the user to ensure that any proprietary rights and existing laws and legislation are observed. © Copyright BASF 2012.  
© AMDRO is a registered trademark licensed to BASF.