


**POISON****KEEP OUT OF REACH OF CHILDREN****READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **DIAZINON 800**

## **INSECTICIDE**

Active Constituent: 800g/L DIAZINON (AN ANTICHOLINESTERASE COMPOUND)

<b>GROUP</b>	<b>1B</b>	<b>INSECTICIDE</b>
--------------	-----------	--------------------

For the control of many insect pests in pastures,  
field crops, vegetables, plantation and orchard crops,  
and commercial and industrial buildings  
(as per the directions for use table)

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.**

A & C RURAL PTY LTD  
A.C.N. 010 865 317  
Lot 96, Potassium Street  
Narangba Qld 4504  
Ph: (07) 3203 3944

**Batch no:**

**CONTENTS: \*\***

**D.O.M.:**

**NRA Approval no: 41698/....**

\*\*This label is to be used for 5Lt, 20Lt pack sizes

**GENERAL INSTRUCTIONS:**

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions, First Aid Instructions and attached leaflet.

**GROUP 1B INSECTICIDE****INSECTICIDE RESISTANCE WARNING**

For insecticide resistance management Country Diazinon 800 Insecticide is a Group 1B insecticide.

Some naturally occurring insect biotypes resistant to Country Diazinon 800 Insecticide and other Group 1B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Country Diazinon 800 Insecticide or other Group 1B insecticides are used repeatedly. The effectiveness of Country Diazinon 800 Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, A & C Rural Pty Ltd accepts no liability for any losses that may result from the failure of Country Diazinon 800 Insecticide to control resistant insects.

Country Diazinon 800 Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, A & C Rural Pty Ltd representative or local agricultural department agronomist.

**COMPATIBILITY:**

This product may be mixed with any one of the following:- Copper Oxychloride, Zineb 800, Wettable Sulphur, Mancozeb, Urea, Winter and Summer Oils.

**MIXING:**

Measure out the required amount of Country Diazinon 800 and add directly to the spray vat with agitation.

**PRECAUTIONS:**

Avoid spray drift onto exposed food, food utensils or food processing machinery. Ventilate treated rooms or buildings before re-occupying.

**SAFETY DIRECTIONS:**

Product is poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing or using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves and full-face respirator. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash contaminated clothing, gloves and respirator. Obtain an emergency supply of Atropine tablets (0.6mg).

**FIRST AID:**

If poisoning occurs, contact a Doctor or Poisons Information Centre (Ph: 13 11 26). IF SWALLOWED, give one Atropine tablet every 5 minutes until dryness of the mouth occurs. IF POISONED BY SKIN ABSORPTION OR THROUGH LUNGS, remove any contaminated clothing, wash skin thoroughly and give Atropine tablets as above. Get to a Doctor or Hospital quickly. Additional information is listed in the Material Safety Data Sheet which is available from A & C Rural Pty Ltd upon request.

**PROTECTION OF LIVESTOCK:**

Dangerous to bees. DO NOT spray any plants in flower while bees are actively foraging. DO NOT spray near fowls, ducks, geese or turkeys. The product is suitable for spraying fowl houses provided birds are removed while spraying is carried out. Avoid spraying drinking water and feed troughs. DO NOT feed grass clippings from treated area to poultry and animals.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT:**

Extremely dangerous to fish. DO NOT contaminate ponds, waterways and drains with this chemical or used container.

**STORAGE AND DISPOSAL AND PROTECTION OF OTHERS:**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**CONDITIONS OF SALE:**

The use of Country Diazinon 800 insecticide being beyond the control of the manufacturer, no warranty expressed or implied is given by A & C Rural Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and A & C Rural Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

**WITHHOLDING PERIOD:**

Do not harvest, graze or cut for stock food crops or pastures for 14 days after application. Do not apply to edible crops within 14 days from harvesting.

**IN A TRANSPORT EMERGENCY,  
DIAL 000,  
POLICE OR FIRE BRIGADE**



**UN 3018  
PESTICIDES, ORGANOPHOSPHORUS LIQUID, TOXIC, N.O.S.**

**EXPIRY**

**DATE:**

**POISON**

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## DIAZINON 800 INSECTICIDE

Active Constituent: 800g/L DIAZINON  
(An Anticholinesterase compound)

**GROUP 1B INSECTICIDE**

For the control of many insect pests in pastures, field crops,  
vegetables, plantation and orchard crops  
(As per the directions for use table)

NRA Approval no: 41698/....

THIS LEAFLET IS PART OF THE LABEL FOR THE 5LT & 20LT  
CONTAINER

A & C Rural Pty Ltd  
Lot 96 Potassium Street  
NARANGBA QLD 4504  
Ph: (07) 3203 3944

**GENERAL INSTRUCTIONS:**

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions, First Aid Instructions and attached leaflet.

**GROUP 1B INSECTICIDE**

**INSECTICIDE RESISTANCE WARNING**

For insecticide resistance management Country Diazinon 800 Insecticide is a Group 1B insecticide.

Some naturally occurring insect biotypes resistant to Country Diazinon 800 Insecticide and other Group 1B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Country Diazinon 800 Insecticide or other Group 1B insecticides are used

repeatedly. The effectiveness of Country Diazinon 800 Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, A & C Rural Pty Ltd accepts no liability for any losses that may result from the failure of Country Diazinon 800 Insecticide to control resistant insects.

Country Diazinon 800 Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, A & C Rural Pty Ltd representative or local agricultural department agronomist.

**COMPATIBILITY:**

This product may be mixed with any one of the following:- Copper Oxychloride, Zineb 800, Wetttable Sulphur, Mancozeb, Urea, Winter and Summer Oils.

**MIXING:**

Measure out the required amount of Country Diazinon 800 and add directly to the spray vat with agitation.

**PRECAUTIONS:**

Avoid spray drift onto exposed food, food utensils or food processing machinery. Ventilate treated rooms or buildings before re-occupying.

**SAFETY DIRECTIONS:**

Product is poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing or using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves and full-face respirator. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash contaminated clothing, gloves and respirator. Obtain an emergency supply of Atropine tablets (0.6mg).

**FIRST AID:**

If poisoning occurs, contact a Doctor or Poisons Information Centre(Ph: 13 11 26). IF SWALLOWED, give one Atropine tablet every 5 minutes until dryness of the mouth occurs. IF POISONED BY SKIN ABSORPTION OR THROUGH LUNGS, remove any contaminated clothing, wash skin thoroughly and give Atropine tablets as above. Get to a Doctor or Hospital quickly. Additional information is listed in the Material Safety Data Sheet which is available from A & C Rural Pty Ltd upon request.

**PROTECTION OF LIVESTOCK:**

Dangerous to bees. DO NOT spray any plants in flower while bees are actively foraging. DO NOT spray near fowls, ducks, geese or turkeys. The product is suitable for spraying fowl houses provided birds are removed while spraying is carried out. Avoid spraying drinking water and feed troughs. DO NOT feed grass clippings from treated area to poultry and animals.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT:**

Extremely dangerous to fish. DO NOT contaminate ponds, waterways and drains with this chemical or used container.

**STORAGE AND DISPOSAL AND PROTECTION OF OTHERS:**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**CONDITIONS OF SALE:**

The use of Country Diazinon 800 insecticide being beyond the control of the manufacturer, no warranty expressed or implied is given by A & C Rural Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and A & C Rural Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

**WITHHOLDING PERIOD:**

Do not harvest, graze or cut for stock food crops or pastures for 14 days after application.  
Do not apply to edible crops within 14 days from harvesting.

**DIRECTIONS FOR USE****COMMERCIAL AND INDUSTRIAL BUILDINGS**

Situation	Pest	State	Rates of Application			Critical Comments
			Sprayer	Mister	Swingfog	
Commercial and Industrial Buildings, Ships, Farm buildings including kennels, stables and piggeries, Refuse areas, Garbage containers	Cockroaches, Silverfish	All States			60mL per litre of fogging oil or distillate	<ul style="list-style-type: none"> <li>DO NOT USE where there is exposed food, food utensils or food processing machinery.</li> <li>DO NOT USE in food preparation areas, e.g. kitchens, tea rooms.</li> <li>Avoid fogging drinking water or feed stuffs</li> </ul>
	Carpet beetles					
	Fleas					
	Flies					
Homes, Flats, Hotels, Bakeries, Canteens, Commercial and Industrial Buildings, Ships, Farm Buildings, Kennels, Stables, Piggeries, Refuse areas, Garbage containers	Cockroaches, Silverfish		6mL per litre of water or kerosene	15mL per litre of water or kerosene		Apply to crevices, cracks, and hiding places, e.g. beneath cupboards, behind sinks and stoves.
	Carpet beetles					Apply to floors and under carpets
	Bed bugs		Spray should be applied under beds, along skirting boards and anywhere bed bugs may be able to shelter, as well as lightly to bedding			
	Fleas		Areas generally infested with fleas should be sprayed as well as sleeping and resting areas of pets. Change infested bedding.			
	Flies		Apply to surfaces on which insects congregate e.g. ceilings, under eaves, walls.			
	Spiders		Remove existing webbing and saturate area with the mixture			
Ants	Apply to ant trails. Attempt to locate nests and thoroughly saturate surface. Use at least 1 litre of mixture per 10 square metres.					
Refuse Areas, Garbage containers	Maggots		60mL per 100 litres of water	15mL per litre of water or kerosene	60mL per litre of fogging oil or distillate	Apply to thoroughly penetrate the refuse.
Ponds, stagnant water	Mosquito larvae		125mL per 100 litres of water	2mL of diesel oil/kerosene per 100 square metres	180mL in fogging oil or distillate per hectare	Apply to breeding area.
Skins and hides	Skin and hide beetles		5mL per litre of water			Apply 60mL of mixture per hide individually, and spray area with 5 litres of mixture per 100 square metres.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## FIELD CROPS AND PASTURES

Crop	Pest	State	Rate/ha Diluted in water. Boom Spray, Aircraft, Mister	Critical Comments
Chou Moellier, Rape, Turnips	Cabbage Moth, Cabbage White butterfly	SA, Vic, WA only	700mL	Spray as necessary. Do not apply when bees are foraging.
Cotton	Cotton flea beetle Red shouldered beetle	NSW, WA	350 to 500mL	Spray when insects are present.
Lucerne	Lucerne Jassid	Qld, WA	350mL	Spray with good coverage when pests are active.
	Spotted Alfalfa aphid	Vic, WA, Only NSW only	700mL 500mL	Spray when there are 20 to 40 aphids/stem on mature plants or 1 to 2 aphids/stem on seedlings. Spray as soon as practical following cutting and repeat as necessary.
Nursery Plants	Aphids, Thrips, Mealy bugs, Scale insects, Plant bugs, Beetles	NSW, Vic, WA	Dripping mixture: 60mL per 100 litres	Thoroughly drench plant material with the dripping mixture. Treatment can be used for plants being transported from NSW into Victoria
Pastures, Cereals (including Maize, Sorghum), Oil Seed crops (including cotton) Pastures, Sugar Cane, Soybeans	Australian Plague Locust	Qld, NSW, SA, Vic, WA only	700mL	Spray directly onto hopper brands and flying swarms.
	Spur Throated Locust, Migratory Locust	Qld, NSW, WA only	700 to 850mL	Spray swarms in early morning or late evening when pests are resting. Preferably spray when locusts are at young stage. Use higher rate when crop is dense or locusts at adult stage.
Pastures	Pasture Webworm	Qld, WA only	700mL	Apply immediately damage is observed – July/November. Spray late in day. Good coverage at ground level is important. Grazing or mowing assists.
Pastures, Cereal Crops	Armyworm, Cutworm	All States	1L	Apply in at least 110Lt water for boom, 22L for aircraft, 22 to 44L for mister. Apply when pests first noticed. (Treatment more effective at very early stage of pest life cycle). Respray may be required within 10 days.
Pastures, Lawns, around trees	Argentine Ants	W.A. only	High Volume: 600mL per 100 litres. Use at least 1 litre per 10 square metres. Spray lawns on grid pattern on one metre squares	Spray thoroughly, ant nests and infested areas. Apply coarse spray to pests and all trees alongside foundation walls, fences, paths and garden beds.
Pasture, Sorghum	Grasshoppers	Vic, Tas, WA only	1 to 1.4L	Spray swarms in early morning or late evening when pests are resting.
Sorghum	Sorghum Midge	Qld, WA only	350mL	Spray from 90% sorghum head emergence to advanced flowering stage at 4 day intervals or as necessary, depending on pest numbers and crop potential. In NSW use 350mL/ha for light infestation and 700mL/ha for heavy infestation or where reinfestation is likely to occur rapidly.
		NSW only	350mL-700mL	
Rice	Bloodworm	NSW, WA only	75 to 150mL	Spray first at or within 24 hours of sowing, by aircraft or ground application equipment to flooded bays. When the water depth is 150mm or less use 75mL/ha, when the water depth is between 150mm and 300mm or when high levels of recently incorporated organic matter are present, use 150mL/ha. Apply further treatments as required until plants are established with satisfactory secondary root development.
	Brown Plant Hopper	Qld, WA only	350mL	Spray when pests are active. For aerial application, use at least 50L spray/ha.

## PLANTATION AND ORCHARD CROPS

Crop	Pest	State	Rate High Volume Per 100L water	Critical Comments
Apples, Pears (dormant)	San Jose Scale	NSW, Vic, SA, WA only	65mL+	Plus add 3L winter oil per 100L. Spray thoroughly two weeks before green tip.
Apples, Pears		Vic, WA only	65mL+	Plus add 1.2L summer oil per 100L. Spray at petal fall, in early December and again in mid-January, if scale serious.
Apples	San Jose Scale, Woolly Aphid (Apples only)	SA, NSW, WA	65mL	Spray at petal fall in early December and again in mid-January, if scale serious. These applications will keep Woolly Aphid in check during the period they are applied. This product will not adversely affect populations of predatory <i>Typhlodromus occidentalis</i> .
			65mL+	Plus add 1.2L summer oil per 100 litres. Apply in autumn if scale or aphid evident in harvest crop.
	Greedy Scale	WA only	65mL+	Plus add 3L winter per 100L. Spray when buds are swelling as a delayed dormant treatment, with thorough coverage of limbs.
			65mL	Spray as required – November to March
Bananas	Banana Weevil Borer	NSW, WA only	125mL	Spray in spring and repeat last summer when adult beetles are present. Apply 600mL of mixture to base of each plant.
	Banana Rust Thrip	Qld, WA only	50mL	Ensure good coverage of fingers and penetration inside bunch. Apply as required when the bunch is fully emerged until bracts have fallen. Apply every 14 days between November and March in North Queensland when Rust Thrips are most active.
Blueberries	Scale Insects	NSW, WA only	70mL	Spray in late afternoon after bees have finished foraging.
Citrus	Spined Citrus Bug	NSW, WA only	65mL+	Thoroughly wet bugs. Do not spray fruit less than 2.5cm diameter to avoid possible blemish. Plus add 1L summer oil per 100L spray.
	Citrus Leaf Miner	Qld, NSW, WA only	65mL	Spray young growth every 10 days while pests active.
	Grasshoppers	Qld, WA only	65mL	Spray when necessary.
Grape Vines	Mealy Bug	Vic, SA, WA only	30mL	Spray during late November-December. For late table grapes, spray early November, again late December, and follow-up if necessary. Thorough coverage is essential. Spray thoroughly as required.
	Australian Plague Locust	WA only	65mL	
Hops	Common and Southern Armyworm	Vic, WA only	65mL	Spray when pests are first noticed. Repeat in 10 days if necessary.
Kiwifruit	Latania Scale, Greedy Scale	NSW, WA	70mL	Apply when the crawlers are leaving the parent scale. A follow-up application may be necessary.
	Leafroller Caterpillar, Cluster Caterpillar			Apply at 10 to 14 day intervals while the caterpillars are active.
Macadamia Nuts	Macadamia Felted Coccid	Qld, NSW, WA	125mL	Spray thoroughly when pests noticed. Repeat applications monthly as necessary.
	Macadamia Leaf Miner	NSW, WA	65mL	Spray in the Spring & autumn when new flush growth is attacked; or as required. Thorough coverage of the outside canopy is required.
Pineapples	Pineapple Scale	Qld, NSW, WA	65mL+	Plus add suitable wetter. Spray plant thoroughly when scales or eggs evident, apply 2000 to 3000L mixture per hectare. Repeat applications at 2-3 week intervals as necessary.
			65mL+	Plus add 1L summer oil per 100L and dip any suspected infected planting material.
	Mealy Bug		Boom spray/ha: 1.5 to 3L	Spray thoroughly as necessary. Use higher rate for heavy infestations.
Stone Fruit	San Jose Scale	SA, WA, NSW, Vic	65mL+	Plus add 3L winter oil/100L spray. Spray thoroughly during dormant period up to budswell
	Green Peach Aphid, Black Cherry Aphid	NSW, WA, SA	65mL	Apply to thorough coverage at budswell. Respray if aphid or scale number build up.
	San Jose Scale	WA, SA, NSW, Vic		

## VEGETABLES

Crop	Pest	State	Rate/Dilution in Water			Critical Comments
			Boom spray/ha	High Volume/ 100L	Per 15L knapsack	
Beans	Bean Fly	Qld, NSW, Vic, WA only	150mL	30mL	-	Spray thoroughly 3, 7 and 14 days after first bean plants emerge. Further weekly sprays may be necessary in heavy attacks.
	Bean Caterpillar		700mL	-	-	Spray when necessary. Some control of aphids will be achieved.
	Blossom Thrips	NSW, Vic, Qld, SA, WA only	350mL	30mL	5mL	Spray when crop is flowering and thrips are in damaging numbers. Repeat as required.
	Seed Maggot	Vic, SA, WA, NSW	3 to 5L	-	-	Spray soil before sowing and work in to a depth of 5 to 8 cm.
Beetroot, Silverbeet, Globe Artichoke	Webworm	Vic, Qld, SA, WA, NSW	700mL	-	-	Spray as necessary. Use 1L/ha for advanced crops. Some control of aphids will be achieved.
Cabbage, Cauliflower, Broccoli, Brussel Sprouts, Kale and Kohlrabi	Cabbage White Butterfly, Centre Grub Cluster caterpillar, Cabbage Aphid, Green Peach Aphid	Qld, WA only	700mL	140mL	30mL	Spray to run off with good coverage. Apply at 10 to 14 day intervals.
	Cabbage White Butterfly, Cabbage Moth	NSW, SA, WA only	700mL to 1.4L	140mL	30mL	Spray to run off with good coverage. Apply at 10 to 14 day intervals. Vary boom spray rate according to plant size. Some control of aphids will be achieved.
	Cabbage White butterfly	Vic only				
	Looper (Chrysodeixis)	Vic, Qld, SA, WA, NSW	700mL	65mL	15mL	
Capsicum, Eggplant	Cutworms	Vic, Qld, SA, WA, NSW	700mL to 1.4L	-	-	
Curcubits	Thrips	Vic, Qld, SA, WA, NSW	350mL	30mL	5mL	Spray when crop is flowering and thrips are in damaging numbers. Repeat as required.
Mushrooms	Mushroom Pests	All States	-	-	-	AT SPAWNING: 140mL in 10L water/1 tonne of moist compost. Mix thoroughly with compost at spawning. AFTER CASING: 30 mL in 10L water/tonne of moist compost. Spray over top of casing soil immediately after casing.
Onions, Garlic	Onion Thrips	NSW, SA, Qld, Vic, WA only	700mL	65mL	-	Spray when thrips are in damaging numbers. Repeat every 10 days as necessary.
Onions	Onion Seedling Maggot	NSW, SA, Vic, WA only	700mL	65mL	-	Spray foliage when flies appear, usually as plants emerge. Respray every 10 days if necessary.
	Onion Maggot, Wireworm	Vic, SA, WA, NSW	3 to 5 L	-	-	Spray soil before sowing and harrow to depth of 5 to 8 cm or irrigate immediately after application.
Potatoes	Potato Moth	Tas, SA, WA only	1L	-	-	Spray immediately 'blister' stage is noticed. Apply at least 170L/ha. In SA, follow up every 14 days as required.
Tomatoes	Thrips	SA, Vic, Qld, WA	350mL	30mL	5mL	Spray from the seedling stage and repeat application at 7-14 day intervals as long as thrips are present. Effective thrips control is necessary to prevent Spotted Wilt virus infection.
	Cutworms	NSW, Vic, Qld, SA, WA	700mL to 1.4L			



22/11/01

COUNTRY DIAZINON 800 INSECTICIDE

PAGE 9 OF 9

Sundry Vegetables (Cantaloupe, Carrots, Celery, Chokos, Cucumbers, Gherkins, Lettuce, Marrows, Pumpkin, Parsnip, Peas, Squash, Watermelons, Rhubarb, Sweet Corn	Caterpillars, Cutworms	NSW, Vic, Qld, SA, WA	700mL to 1.4L			Spray as necessary. Vary rate according to plant size. Some control of aphids and thrips will be achieved.
--	------------------------	-----------------------------	------------------	--	--	--

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:** Do not harvest, graze or cut for stock food crops or pastures for 14 days after application.  
Do not apply to edible crops within 14 days of harvesting.

**UN 3018  
PESTICIDES, ORGANOPHOSPHOROUS, LIQUID, TOXIC, N.O.S.**

**IN A TRANSPORT EMERGENCY,  
DIAL 000,  
POLICE OR FIRE BRIGADE**

