

Main panel

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

 **Temprid® 75**

RESIDUAL INSECTICIDE

**ACTIVE CONSTITUENTS: 50 g/L IMIDACLOPRID
 25 g/L BETA-CYFLUTHRIN**

GROUP 3A 4A INSECTICIDE

1 L
5 L

A residual insecticide concentrate for the control of various insect pests in domestic, commercial, industrial and public buildings and facilities as per the Directions for Use.

IMPORTANT: Read the attached booklet before use.



Booklet front page

CAUTION

KEEP OUT REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TEMPRID® 75 RESIDUAL INSECTICIDE

ACTIVE CONSTITUENTS: 50 g/L IMIDACLOPRID

25 g/L BETA-CYFLUTHRIN

A residual insecticide concentrate for the control of various insect pests in domestic, commercial, industrial and public buildings and facilities as per the Directions for Use.

STORAGE AND DISPOSAL

Store in tightly closed, original container in a safe, well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight.

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the skin and eyes. Avoid contact with skin and eyes. When preparing product for use and using the prepared spray, wear cotton overalls over normal clothing buttoned to the neck and wrist and elbow length chemical-resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766. Remove from contaminated area. Apply artificial respiration if not breathing.

SAFETY DATA SHEET

Additional Information is listed in the Safety Data Sheet which may be found at www.bayeres.com.au.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

IMPORTANT: READ THIS BOOKLET BEFORE USE.

APVMA Approval Number: 64371/58117

Temprid® is a Registered Trademark of the Bayer Group.

Bayer CropScience Pty. Ltd.

A.B.N. 87 000 226 022

391-393 Tooronga Rd

Hawthorn East, Vic. 3123

Phone: (03) 9248 6888

Fax: (03) 9248 6803

Technical enquiries: 1800 804 479

Website: www.bayeres.com.au



(For 5 L containers only)

For 24 hour specialist advice in an emergency only phone 1800 033 111

Batch Number:

DOM:

Booklet inside

DIRECTIONS FOR USE

RESTRAINT

DO NOT apply spray treatment outdoors if heavy rains are expected to occur within 48 hours of application, or if area is about to be irrigated.

SITUATION	PEST	RATE	CRITICAL COMMENTS
Domestic, commercial, industrial and public buildings and facilities (including but not limited to food processing establishments, shops, factories, farm buildings, ships, trains, aeroplanes, offices, schools, storerooms, hospitals, barracks, private houses and external areas associated with such establishments)		Non-porous surfaces: 50 ml / up to 5 L water / 100 m ² Porous surfaces or through use of power equipment: 50 ml / up to 10 L water / 100 m ²	<p>Best-Practice Recommendation:</p> <ul style="list-style-type: none"> • Estimate area of surface which needs to be treated. • Calibrate spray equipment to identify volume of spray necessary to achieve coverage of target area. • Measure out required volume of Temprid® 75 Residual Insecticide and mix according to General Instructions below. • DO NOT exceed point of run-off. • Non-porous surfaces: Apply as a coarse spray.
	Ants		<p>DIRECT NEST TREATMENT: Where possible locate the nest and spray directly into nest entrances. Penetration of insecticide can be assisted by pre-wetting sandy and porous soils with water. A thorough soaking treatment with insecticide mixture is required to destroy the nest.</p> <p>TREATMENT WHEN NESTS CANNOT BE LOCATED: Apply to all ant trails and areas where ants are foraging. Search for ants underneath materials (e.g. rocks, timber) and apply to these areas. Where indoor infestation has been identified it is recommended that an external perimeter treatment be installed (refer below) and that indoor areas are treated with a suitably registered bait product such as from the Maxforce® ant bait range.</p> <p>EXTERNAL PERIMETER TREATMENTS: Apply to all potential entrance points of the structure (e.g. cracks and crevices in brickwork and gaps in timber weatherboards). Treat the outside of the structure; walls, and around doors, windows, and other areas where ants are active or may enter or hide. Ensure application in wall voids or other enclosed cavities is made with spray nozzle adjusted to produce a fine mist in order to avoid run-off. Treat the soil, turf or other ground covering adjacent to the structure (in a band up to 3 metres wide) where ants are seen, have been found, or can find shelter.</p>

SITUATION	PEST	RATE	CRITICAL COMMENTS
Domestic, commercial, industrial and public buildings and facilities (including but not limited to food processing establishments, shops, factories, farm buildings, ships, trains, aeroplanes, offices, schools, storerooms, hospitals, barracks, private houses and external areas associated with such establishments)	Cockroaches	Non-porous surfaces: 50 ml / up to 5 L water / 100 m ² Porous surfaces or through use of power equipment: 50 ml / up to 10 L water / 100 m ²	Conduct a thorough inspection and apply to known and potential cockroach harbourages. A barrier application around areas where cockroach infestations may be originating is also suggested. NOTE: The use of an Insect Growth Regulator (such as Starycide® Insect Growth Regulator) is considered a useful supplementary treatment for cockroach control when cockroach infestation is severe. Repeat application as required. Where application of a spray is not suitable it is recommended that a suitably registered gel such as from the Maxforce® Gel range be applied as per the registered label on that product.
	Crickets		Spray infested areas. In addition a barrier treatment can be sprayed across areas where these pests enter buildings.
	Mosquitoes and flies		Spray surfaces where flies and mosquitoes are known to come to rest. Flies – include external surfaces that are exposed to the morning sun. Mosquitoes – spray walls, awnings, fences and vegetated areas in proximity to buildings. DO NOT apply to edible plants. NOTE: The perception of the level of control of flying insects achieved within any given area treated with a residual surface spray will be affected by the extent of in-migration of individuals from external (untreated) areas. A residual surface spray will only control those insects that alight on the treated surfaces and therefore complete elimination of flying insects from the treated area should not necessarily always be expected.
	Millipedes		Apply as a coarse spray to infested areas. Recommended for use on non-porous surfaces only.
	Paper nest wasps		Apply diluted spray to the point of run-off directly to the papernest ensuring thorough and even coverage. When all adult wasps have been knocked down the nest may be safely removed from the structure.
	Spiders		Apply to areas where spiders are known to occur, their webbing and areas where spiders may hide. It is recommended that any webbing is left in place for a few days after application to optimise efficacy.
Internal areas and surrounds of domestic, commercial, industrial and public buildings and facilities	Fleas		Clean and vacuum infested areas and dispose of dust collected in the garbage in a sealed plastic bag. Lightly spray infested areas concentrating on areas under furniture etc. NOTE: The use of an Insect Growth Regulator (such as Starycide® Insect Growth Regulator) is considered a useful supplementary treatment for flea control. Treat pets with a suitably registered flea control product. See also the note on application to carpets below.

SITUATION	PEST	RATE	CRITICAL COMMENTS
Domestic, commercial, industrial and public buildings and facilities (including but not limited to hotels and backpacker accommodation, private dwellings, barracks, hospitals, ships, trains, aeroplanes, offices, schools)	Bedbugs	Non-porous surfaces: 50 ml / up to 5 L water / 100 m ² Porous surfaces or through use of power equipment: 50 ml / up to 10 L water / 100 m ²	Professional Pest Managers undertaking a bedbug treatment are expected to be familiar with the AEPMA Code of Practice (CoP) for the Control of Bedbug Infestations in Australia and take into account the recommendations included within this document in relation to bedbug management. Prior to treatment, a thorough inspection of all infested and potentially infested areas should be carried out. Treatment should be directed to all areas of infestation, e.g. the bed-frame and walls, cracks, crevices and skirting boards in the vicinity of the bed. It may also be applied to the edges and seams of the mattress but DO NOT treat the flat surfaces which people sleep on. Consideration should also be given to appropriate non-chemical methods of control in areas where insecticide application is not possible. Control of pyrethroid-resistant strains of bedbug is dependent on direct spray application rather than residual activity. Re-inspection is essential (in accordance with the recommendations of the CoP).
	Carpet beetles		Apply to infested carpets particularly in areas of little or no traffic and underneath windows where adult beetles may be entering the building. See also the note on application to carpets below.
	Clothes moths		Spray surfaces in areas where clothes are stored. DO NOT spray the clothes themselves.
	Silverfish		Direct spray to areas of potential and existing infestations e.g. under skirting boards and furniture where dark conditions may favour silverfish populations. If spraying inside cupboards or storage areas remove clothing and utensils etc. before spraying. NOTE: The use of an Insect Growth Regulator (such as Starycide® Insect Growth Regulator) is recommended as a supplementary treatment for silverfish control.
	Skin and hide beetles		Apply as a coarse spray paying particular attention to the flesh side of the hides.
	Stored product pests		The use of this product in the control of stored product pests should be considered part of an integrated strategy to prevent development of insect populations. Best results will be obtained where good hygiene is maintained. Infested areas should be cleaned prior to application and regularly cleaned thereafter to reduce food sources for pests. When treating for beetle pests pay special attention to sheltered areas such as cracks and crevices, under furniture and equipment, floors, walls and shelves. Adult moths can be controlled by applying this product to walls and surfaces where the flying adult moths come to rest. DO NOT apply to food or onto food preparation areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Mixing:

Add the required quantity of **Temprid® 75 Residual Insecticide** to water while stirring.

Application:

DO NOT spray the moving parts of any machinery, electric motors or switchgear with the water-based product.

Compatibility:

Temprid® 75 Residual Insecticide is physically compatible with **Starycide® Insect Growth Regulator**.

INSECTICIDE RESISTANCE WARNING

GROUP	3A	4A	INSECTICIDE
-------	----	----	-------------

For insecticide resistance management **Temprid® 75** is a Group 3A & 4A insecticide. Some naturally occurring insect biotypes resistant to **Temprid® 75** and other Group 3A and 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if **Temprid® 75** or other Group 3A and 4A insecticides are used repeatedly. The effectiveness of **Temprid® 75** on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty. Ltd. accepts no liability for any losses that may result from failure of **Temprid® 75** to control resistant insects.

For further information on managing insecticide resistance contact your local supplier or Bayer Environmental Science representative.

PRECAUTIONS

DO NOT apply using spraying equipment carried on the back of the user.

For surface treatment only – DO NOT spray into the atmosphere using foggers or other devices designed to create an aerosol.

When using in animal houses DO NOT spray directly onto livestock, their food, water, litter, feed bins, or surfaces that are accessible to the stock.

DO NOT spray directly on humans, exposed food or food utensils or food preparation areas.

DO NOT spray clothing.

Avoid contact of spray with timber to which a stain is subsequently to be applied, unless the owner is notified that the timber will need additional preparation, consistent with that if water alone had contacted the timber. (Evidence of water contact on timber is sometimes only evident once the stain has been applied).

BEFORE USE - Remove or cover all exposed foodstuffs. Cover all dishes and utensils and surfaces where food is prepared.

AFTER USE - Thoroughly ventilate treated areas. Clean up thoroughly before processing/serving resumes.

Note on application to carpets:

Whilst carpets are regarded as porous surfaces, excessive liquid application may produce water-stains and therefore the non-porous surface rate should be adopted if possible. Hence for short pile carpet, the 5 L dilution rate (per 100 m²) should be sufficient. However, for long pile or shag carpet, the 10 L dilution rate (per 100 m²) may be required, to ensure effective penetration and application to all of the carpet-pile. Since carpets vary in their thickness, colour and material it is recommended that a small, inconspicuous area of carpet be sprayed and allowed to dry first to ensure that staining will not occur. Allow treated carpets to dry before resuming activity on them.

RE-ENTRY PERIOD

DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers. Cover fish tanks before using.

STORAGE AND DISPOSAL

Store in tightly closed, original container in a safe, well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging

to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the skin and eyes. Avoid contact with skin and eyes. When preparing product for use and using the prepared spray, wear cotton overalls over normal clothing buttoned to the neck and wrist and elbow length chemical-resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766. Remove from contaminated area. Apply artificial respiration if not breathing.

SAFETY DATA SHEET

Additional Information is listed in the Safety Data Sheet which may be found at www.bayeres.com.au.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty. Ltd. accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval Number: 64371/58117

Temprid® is a Registered Trademark of the Bayer Group.

Bayer CropScience Pty. Ltd.
A.B.N. 87 000 226 022
391-393 Tooronga Rd
Hawthorn East, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6803
Technical enquiries: 1800 804 479
Website: www.bayeres.com.au

For 24 hour specialist advice in an emergency only phone 1800 033 111
