Fendona® Plus 60SC

Insecticide

A high performance residual insecticide providing rapid knockdown and broad spectrum control

General description

Fendona Plus 60SC is the first choice synthetic pyrethroid spray treatment for insect pests. It is a high performance residual insecticide with a broad spectrum of action against a large variety of insect pests. The unique **Fendona Plus** formulation provides superior surface adherence though the formation of micro-crystalline particles.

Fendona Plus is suitable for use in and around domestic households, industrial and commercial establishments including food processing establishments, offices, schools, store rooms, and hospitals. **Fendona Plus** is now registered for use around agricultural buildings.



Features	Benefits Fendona Plus controls a wide range of flying and crawling insects in domestic, commercial, agricultural and public health applications. In particular, it gives highly effective control of cockroaches, ants, bed bugs, silverfish, fleas, flies, mosquitos, meal moths and spiders.		
Broad spectrum control			
Rapid knockdown	Fendona Plus is economic to use and provides superior efficacy when compared to other 3A insecticides. It delivers immediate results with insects showing the signs of sensitisation within 5 minutes, full knockdown within 30 minutes.		
New low dose formulation	The advanced low dose formulation contains 60g/l of alpha-cypermethrin ensuring dependable results at exceptionally low doses for maximum application convenience and minimum total treatment cost.		
Superior residual action The unique micro-crystalline suspension concentrate formulation results in even surface of excellent adherence to porous and other difficult surfaces like mortar, brick and cement we active ingredient absorption. The micro-crystalline formulation also results in the active ingreadily bio-available resulting in superior control of difficult pests such as bed bugs.			

Application

Fendona Plus controls insects by contact action and ingestion. When applied as a surface spray, the product offers long residual activity. **Fendona Plus** insecticide can be applied with conventional spray equipment used to control public health pests.

To obtain best results, all surfaces should be treated, especially cracks, crevices and infested areas. A higher volume of water may be required on very absorbent surfaces. Fill the spray tank with half the volume of required water. Add the required amount of **Fendona Plus** insecticide. Top up the remainder of the water and agitate thoroughly.

Packaging

Now available in 1 Litre bottles



Fendona® Plus 60SC

Insecticide

Areas	Pest	Rate	Critical comments
Domestic, industrial, commercial, public buildings and around agricultural buildings: - including food processing establishments, shops, restaurants factories, ships, offices, schools, store rooms, hospitals, barracks, private houses and external areas associated with such establishments.	Cockroaches, Fleas, Flies, Silverfish, Spiders, Mosquitoes, Bed bugs Indian Meal Moth.	Initial treatment and heavy infestations: 10 mL in 1L water	For crawling insects, apply as a coarse low pressure spray to areas where pests frequent.
			For flying insects, apply as a surface treatment using a fine spray to wet the surface.
			Apply 5L of diluted spray mix/100m2 up to but not exceeding the point of run-off.
		Maintenance treatment and light infestations: 5 mL in 1L water	For cockroach control, where the use of a spray is not suitable, a registered cockroach gel product such as GOLIATH® Gel or AMDRO® Gel may be applied as per the registered label.
			For spider control, spray spiders, webbings and areas where spiders my hide directly. The treated webs should be left in place for at least 72 hours to increase efficacy, then removed for monitoring or any re-infestation.
			For flies and mosquitoes treat surfaces where flies and mosquitoes are observed to rest.
			For bedbugs treat bedframes, behind bed heads, bedside tables, cupboards cracks, crevices, carpet edge, and skirting boards. FENDONA® may also be applied to the edges and seems of the mattress, but do not treat the flat surface on which people sleep. An IPM approach should be followed for the control of bed bugs. For more information refer to the Code Of Practice for the Control of Bed Bug Infestations in Australia.
			For Indian meal moth, following the removal of any food, animal feed and containers, apply to areas where pests are present.
			Add a non-ionic surfactant at the registered label rates to improve spreading on hard to wet surfaces.
	Ants	Initial treatment, extended control and heavy infestations: 16 mL in 1 L water short term control and light infestations: 9 mL in 1 L water	Follow the directions for use as per crawling insect treatment (above).

For technical advice and product information visit www.pestcontrol.basf.com.au

