

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

RESOLV[®]

NO WAX SOFT BAIT

SOFT BAIT =
PALATABLE
AND
ATTRACTIVE



SINGLE
FEED

ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE

Highly palatable soft bait for the control of rats and mice



FOR RATS AND MICE



Contains Bitrex[®], a human taste deterrent

Emergency Phone No: 1800 033 111

**LIPHA
TECH**

— GROUPE DE SANGOSSE

Manufactured by:
LIPHATECH S.A.S.
Bonel BP 3
47480 PONT DU CASSE,
France

Distributed by:
De Sangosse Australia Pty Ltd
38 Ricketty St, Mascot NSW 2020
Ph: 02 9078 7859
infoau@desangosse.com

NET CONTENTS 5.5KG

Contains 550 x 10g baits



DIRECTIONS FOR USE

RESTRAINTS

DO NOT exceed 3 m between bait stations for mice or 9 m for rats in clean-out treatments. DO NOT place baits in open. DO NOT remove bait from sachet.

SITUATION	PESTS	RATE	CRITICAL COMMENTS
In and around agricultural, commercial, domestic and industrial situations	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>)	Normally 1 to 10 soft baits per bait station are sufficient at the first application. For further applications, refer to Critical Comments.	The feeding behaviour of rats and mice is slightly different. Rats feed in large quantities in few locations whereas mice are more erratic and tend to feed in different places at different times. First determine the infested areas. Where droppings are seen, inside ceiling cavities, along runs are locations more probably frequented by rats and mice. Remove as much alternative food as possible. Check bait consumption after 3 days. Replace eaten baits. Where a high level of consumption is noticed, place more baits. Remove baits in new locations where there has been no bait taken.
	Mice (<i>Mus musculus</i>)	Normally 1 to 5 soft baits per bait station are sufficient at the first application. For further applications, refer to Critical Comments.	Inspect again 3 days later and then once a week until feeding ceases. Normally treatment takes 2 to 3 weeks. Set out bait stations in places frequented by rats and mice in and around buildings (within 2 m) or enclosed spaces, e.g. drains. Clean-Out Treatment: For Rats: Place bait stations at intervals of 5 to 9 m in infested areas. For Mice: Place bait stations at intervals of 2 to 3 m in infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Resolv® Soft baits contain a human taste deterrent which prevents accidental consumption by children. Try not to touch the bait as this may deter rodents from eating it. Please use gloves. Usually rodents will die within 3 to 5 days after eating the bait. If reinfestation should occur, you can repeat treatment once. For further treatments, it is strongly advised to alternate with a product containing a different active ingredient to avoid the development of genetic resistance.

BAITING STRATEGY

- Eliminate as far as practical, all alternative food sources.
- Bait infested area. Place baits under cover.
- Actively search for, remove and destroy dead and moribund rodents and rodent faeces whilst bait is being used.
- Ensure baits are removed when rodent activity has ceased.

Clean-Out Treatments (Pulse Baiting)

- Clean-out treatments are used to remove existing infestations.
- Always record the number and locations of bait sachets for the purpose of bait replacement and later removal.
- For rats, place baits at intervals of 5 to 9 m in infested areas.
- For mice, place baits at intervals of 2 to 3 m in infested areas.
- The use of secure tamper resistant bait stations is preferred.
- Place baits so as to prevent access to non-target animals. Where access to non-target animals cannot otherwise be prevented, secured tamper resistant bait stations must be used.
- Use the greater number of baits and shorter distance between bait placements when infestations are heavy.
- Conduct initial inspection of bait points after 3 or 4 days. Replace any bait that has been consumed.
- Repeat approximately 4 days later and then weekly. Replace contaminated or spoiled baits.
- Baiting for at least 2 weeks (3 or 4 pulses) will be necessary to reduce rat/mouse numbers to a low level, although heavy infestations may require longer treatments to achieve complete eradication.
- Discontinue the treatment when effective control has been achieved and remove all bait and bait stations unless a maintenance treatment is to be undertaken.

Monitoring and/or Maintenance Treatments

- Monitoring and maintenance treatments are required at sensitive sites where there is no current rodent infestation but where the prevention of such infestation is essential to prevent significant risks to human health. Such treatment may be a condition for audit approval within food safety guidelines. However, all use of this product must comply with the conditions of this label.
- Always record the number and locations of bait stations for the purpose of bait replacement and later removal.
- Place baits in areas prone to rodent infestation.
- Place baits so as to prevent access to non-target animals. Always use tamper-resistant bait stations for monitoring and maintenance treatments and fix the stations securely to the substrate.
- Bait stations should be checked at regular intervals according to an agreed and documented schedule.
- Use placebo or unpoisoned baits if the placement of the bait away from buildings presents a hazard to non-target animals.
- When a rodent infestation is detected the operator should revert to the directions for Clean-Out treatments.

PRECAUTIONS

The use of this product for the control of protected native rodents requires permission from wildlife authorities. Do not allow bait to contaminate surfaces that may come into contact with food or water intended for human or animal consumption. Do not apply bait to crops. Place baits in locations or bait stations which are inaccessible to children, livestock, other animals and birds. This product and dead or affected rodents are hazardous to pigs, cats, dogs, native wildlife and poultry. This product is restricted to use in and around buildings (within 2 metres) or enclosed spaces (e.g., drains).

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Do not store for prolonged periods in direct sunlight.

Break, crush, puncture, and bury empty containers and dispose of unused bait in an approved waste management facility. If an approved waste management facility is not available, bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Empty containers and product should not be burnt. Do not use containers which have held bait for any other purpose. Dispose of carcasses safely by burning or burying in an approved landfill. Do not place dead rats or mice in refuse bins or rubbish tips. Dispose of spoiled baits by placing in plastic bags and putting in garbage.

SAFETY DIRECTIONS

Poisonous if swallowed. Repeated minor exposure may have a cumulative poisoning effect. When using the product, wear rubber gloves. Wash hands after use. After each day's use, wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26. Vitamin K (phytonadione) is antidotal.

Additional information is listed in the Safety Data Sheet.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

This product as supplied is of a high grade and believed to be suitable for any purpose for which it is expressly recommended and must be used in accordance with the Directions for Use given on the label. No responsibility is accepted in respect of the product save those non-excludable conditions implied by any Federal and State legislation or law of a territory.

Resolv® Registered trademark of LIPHATECH S.A.S.
Bitrex® Registered trademark of Macfarlan Smith Limited

BN

DOM

APVMA Approval No. 86179/114907



LIPHATECH

— GROUPE DE SANGOSSÉ —