

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**Maxforce<sup>®</sup>**

**COCKROACH GEL**

**Active Constituent: 16.5 g/kg HYDRAMETHYLNON**

**A gel bait for the control of cockroaches by professional pest control operators as per the DIRECTIONS FOR USE.**

## **GENERAL INSTRUCTIONS**

### **DIRECTION FOR LOADING RESERVOIRS and OPERATION**

On the handle of the drive rod point the arrow facing up. Pull the drive rod as far as it will go. Attach a pre-filled reservoir of MAXFORCE gel to the bait injector with a twist locking movement. Break off and dispose of the plastic tip and attach the dispensing cap. Push the drive rod until the plunger seats against the piston in the reservoir of the gel. Turn the drive rod knob until teeth are facing down and engaging the trigger mechanism (arrow down). Gently squeeze the trigger until the gel flows out.

### **DIRECTIONS FOR CHANGING RESERVOIRS**

With the arrow on the drive rod pointing up, pull the drive rod out as far as it will go. Remove the empty reservoir, wrap and place in rubbish bin. Reservoirs that still contain gel can be sealed with the supplied cap for further use. To attach a new reservoir, follow the directions above.

Maxforce Cockroach Gel is an insect stomach poison. The bait will adhere to non-greasy and non-dusty surfaces and will remain pliable and palatable to cockroaches as long as it is visibly present.

Re-apply when bait is no longer visibly present, according to the remaining level of infestation. A visible inspection of bait placements is recommended at one month after the initial treatment. Re-application should occur if the bait is visibly depleted. Pest population reduction will be apparent within a week or less. Avoid contact with porous surfaces as the bait may stain.

## **PRECAUTIONS**

Avoid contact with food, food utensils or surfaces where food is prepared or stored. Do not apply on food preparation or serving surfaces. Should not be used in areas accessible to children. Do not use on or around electrical equipment due to the possibility of electric shock. DO NOT apply the product in areas easily accessible to children and animals.

## **STORAGE AND DISPOSAL**

Store in the closed original container in a cool well ventilated area. Do not store for prolonged periods in direct sunlight. Dispose of used tube by wrapping in paper, placing in plastic bag and putting in garbage.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.  
DO NOT apply the product in areas easily accessible to children and animals.

## **SAFETY DIRECTIONS**

Harmful if swallowed. When using the product wear rubber gloves. Wash hands after use. After each days use wash gloves.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Telephone: 13 11 26).

## **MATERIAL SAFETY DATA SHEET**

Additional information is listed in the material safety data sheet.

## **EXCLUSION OF LIABILITY**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Maxforce® is a registered trademark of Bayer

NRA Approval Number: 56626/0203

FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY  
PHONE 1800 033 111

**DIRECTIONS FOR USE**

**Restraint**

DO NOT place in areas that are routinely washed.

**PEST:** For the control of German and other common cockroaches.

**RATE:** Apply as 1-2 spots or in a band 75 to 150 mm long per square metre. Recommended spot size ¼ to ½ g.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**