

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SOREXA[®] Pro

RODENTICIDE BLOCKS

ACTIVE CONSTITUENT: 0.05 g/kg DIFENACOU



SOREX
INTERNATIONAL

READY TO USE BAIT FOR THE CONTROL OF BROWN OR NORWAY RAT (*Rattus norvegicus*) AND MICE, including those resistant to other anticoagulants. FOR USE IN DAMP OR DRY SITUATIONS, in and around domestic, industrial, commercial, public services, agricultural buildings.



net contents

5kg

A PROFESSIONAL PEST CONTROL PRODUCT



LABL01094

Manufactured by: Sorex Limited, St Michael's Industrial Estate, Widnes, Cheshire, England W48 8TJ. Marketed by Sorex International, 31 Dover Street, Red Hill, QLD 4059. Telephone: 0412 251016.



APPROVED
LABEL

File name:	61842_41643_5KG_Label_MPL_V01.pdf
Dimensions:	158 x 200mm
Scale:	90% ON A4
Date:	28th May 2007

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT place blocks in the open.

DO NOT place blocks in locations accessible to children and domestic animals.

PEST	SITUATION	RATE	CRITICAL COMMENTS
Mice	For use in and around (within 2 metres) industrial, commercial, public services, domestic, agricultural buildings (machinery sheds etc).	Use 1 block per bait point	Explore the site and locate runs, nests, droppings and signs of damage. Eliminate as far as practical, all alternative food sources. Place bait points wherever droppings, damage or other signs of activity are seen. As mice are sporadic feeders use many bait points, not more than 3m apart.
Brown or Norway Rat (<i>Rattus norvegicus</i>)		Use up to 5 blocks per bait point	Explore the site and locate runs, burrows, droppings and signs of damage. Eliminate as far as practical, all alternative food sources. Place a number of bait points in dry locations throughout the infested area, on runs, in burrows and where droppings are seen, not more than 9m apart.
General information and baiting strategy for rats and mice: Eliminate as far as practicable, all other food sources. Always place bait where it is inaccessible to children, animals and birds. Protect and cover bait. When using outdoors, it is best practice to use weather resistant bait stations. It is recommended that lockable, securable, weather resistant bait stations be used where possible. Alternatively place bait in suitable containers or use weather resistant bait stations constructed from drainpipes or boxes. Fix, nail or wire blocks in place where appropriate. Try not to touch the bait as this deters mice and rats from eating it. Do not place baits where food, feed or water could become contaminated.			Record the number and locations of bait stations for the purpose of replacement and later removal. Inspect baits frequently, particularly in the first ten days. Replace eaten bait. If bait is completely consumed, the quantity should be increased. Continue replenishing bait until left untouched. Baiting for at least 2 weeks will be necessary to reduce rat/mouse numbers. Most deaths occur 4 to 7 days after consumption of bait, but deaths may occur for up to 2 weeks. During treatment search for and collect dead rats or mice. Following treatment remove all remains of bait and bait containers and search for and collect dead rats or mice. Burn or bury carcasses. Do not place in refuse bins. DO NOT let cats, dogs or non-target animals eat poisoned mice and rats. While handling remaining bait material/animal carcasses, wear rubber gloves.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**GENERAL INSTRUCTIONS**

Sorex[®] Pro Rodenticide Blocks is a palatable all-purpose rodenticide formulation based on the advanced anticoagulant, difenacoum, which is equally effective against rats and mice, including strains resistant to some other anticoagulants. In some circumstances more than one feed may be required to produce mortality that generally occurs 4-10 days after consuming the bait.

Sorex[®] Pro Rodenticide Blocks are especially suited for use outdoors and in damp situations and are ideal for burrow baiting. The central hole in the block allows for fixing in bait stations, and for nailing or wiring in place to help prevent block carriage by rodents. Indoors, Sorex[®] Pro Rodenticide Blocks can be used in homes, factories, public buildings industrial premises and offices, as well as hotels, restaurants and food storage facilities where bait security is important. Sorex[®] Pro Rodenticide Blocks contains Bitrex[®] (a human taste deterrent) to help prevent accidental human consumption.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

The bait is hazardous to pigs, cats, dogs, poultry, birds, and wildlife and care should be exercised in its use. DO NOT place baits in locations that are accessible to domestic animals, livestock, birds or other animals. DO NOT contaminate dams, waterways and drains with bait or its used containers. DO NOT use this product to control native animals. The product is restricted to use in and around buildings (within 2 metres) or within enclosed spaces. In areas where possums and other non-target animals frequent, baits should be secured inside proper bait stations or bait stations constructed from drainpipes or boxes.

STORAGE AND DISPOSAL

Store in original container, tightly closed, in a safe place. Triple or preferably pressure rinse containers before disposal. Dispose of rinsings and unused or untaken bait in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit. Empty containers and product should not be burnt. Search for dead mice and rats, burn or bury them, do not place in refuse bins or rubbish tips. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased. Containers that have held bait should not be used for any other purpose.

SAFETY DIRECTIONS

Poisonous if swallowed. Will irritate the skin. May irritate the eyes. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. If

on skin and after each baiting, wash thoroughly with soap and water. Use scoop or measure. Vitamin K1 (phytomenadione) is antidotal. When opening the container and using the product, wear elbow-length rubber gloves. After each day's use, wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26. Vitamin K1 (phytomenadione) is antidotal.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

NOTE TO PHYSICIANS AND VETERINARIANS

Difenacoum, the active constituent of Sorex[®] Pro Rodenticide Blocks, is a long lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

This product as supplied is of a high grade and believed to be suitable for any purpose for which it is expressly recommended and must be used in accordance with the Directions for Use given on the label. No responsibility is accepted in respect of the product save those non-excludable conditions implied by any Federal and State legislation or law of a Territory.

APVMA Approval No. 61842/5KG/0407

Batch No.

® Sorex[®] is a registered trademark of Sorex Limited.

D.O.M.

® Bitrex[®] is a registered trademark of Macfarlan Smith Limited.

LABL01095



SOREXA[®] Pro
RODENTICIDE BLOCKS



File name:	61842_41643_5KG_Label_MPL_V01.pdf
Dimensions:	158 x 200mm
Scale:	90% ON A4
Date:	28th May 2007

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SOREXA[®] Pro

RODENTICIDE BLOCKS

ACTIVE CONSTITUENT: 0.05 g/kg DIFENACOUM



**SOREX
INTERNATIONAL**

net contents

10KG

READY TO USE BAIT FOR THE CONTROL OF BROWN OR NORWAY RAT (*Rattus norvegicus*) AND MICE, including those resistant to other anticoagulants. FOR USE IN DAMP OR DRY SITUATIONS, in and around domestic, industrial, commercial, public services, agricultural buildings.

A PROFESSIONAL PEST CONTROL PRODUCT



Manufactured by: Sorex Limited, St Michael's Industrial Estate, Widnes, Cheshire, England WA8 8TJ. Marketed by Sorex International, 31 Dover Street, Red Hill, QLD 4059. Telephone: 0412 251016.

LABL01096



**APPROVED
LABEL**

File name:	61842_41643_10KG_Label_MPL_V01.pdf
Dimensions:	341mm x 179mm
Scale:	70% on A4
Date:	28th May 2007

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT place blocks in the open.

DO NOT place blocks in locations accessible to children and domestic animals.

PEST	SITUATION	RATE	CRITICAL COMMENTS
Mice	For use in and around (within 2 metres) industrial, commercial, public services, domestic, agricultural buildings (machinery sheds etc).	Use 1 block per bait point	Explore the site and locate runs, nests, droppings and signs of damage. Eliminate as far as practical, all alternative food sources. Place bait points wherever droppings, damage or other signs of activity are seen. As mice are sporadic feeders use many bait points, not more than 3m apart.
Brown or Norway Rat (<i>Rattus norvegicus</i>)		Use up to 5 blocks per bait point	Explore the site and locate runs, burrows, droppings and signs of damage. Eliminate as far as practical, all alternative food sources. Place a number of bait points in dry locations throughout the infested area, on runs, in burrows and where droppings are seen, not more than 9m apart.

General information and baiting strategy for rats and mice: Eliminate as far as practicable, all other food sources. Always place bait where it is inaccessible to children, animals and birds. Protect and cover bait. When using outdoors, it is best practice to use weather resistant bait stations. It is recommended that lockable, securable, weather resistant bait stations be used where possible. Alternatively place bait in suitable containers or use weather resistant bait stations constructed from drainpipes or boxes. Fix, nail or wire blocks in place where appropriate. Try not to touch the bait as this deters mice and rats from eating it. Do not place baits where food, feed or water could become contaminated.

Record the number and locations of bait stations for the purpose of replacement and later removal. Inspect baits frequently, particularly in the first ten days. Replace eaten bait. If bait is completely consumed, the quantity should be increased. Continue replenishing bait until left untouched. Baiting for at least 2 weeks will be necessary to reduce rat/mouse numbers. Most deaths occur 4 to 7 days after consumption of bait, but deaths may occur for up to 2 weeks. During treatment search for and collect dead rats or mice. Following treatment remove all remains of bait and bait containers and search for and collect dead rats or mice. Burn or bury carcasses. Do not place in refuse bins. DO NOT let cats, dogs or non-target animals eat poisoned mice and rats. While handling remaining bait material/animal carcasses, wear rubber gloves.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Sorex[®] Pro Rodenticide Blocks is a palatable all-purpose rodenticide formulation based on the advanced anticoagulant, difenacoum, which is equally effective against rats and mice, including strains resistant to some other anticoagulants. In some circumstances more than one feed may be required to produce mortality that generally occurs 4-10 days after consuming the bait.

Sorex[®] Pro Rodenticide Blocks are especially suited for use outdoors and in damp situations and are ideal for burrow baiting. The central hole in the block allows for fixing in bait stations, and for nailing or wiring in place to help prevent block carriage by rodents. Indoors, Sorex[®] Pro Rodenticide Blocks can be used in homes, factories, public buildings industrial premises and offices, as well as hotels, restaurants and food storage facilities where bait security is important. Sorex[®] Pro Rodenticide Blocks contains Bitrex[®] (a human taste deterrent) to help prevent accidental human consumption.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

The bait is hazardous to pigs, cats, dogs, poultry, birds, and wildlife and care should be exercised in its use. DO NOT place baits in locations that are accessible to domestic animals, livestock, birds or other animals. DO NOT contaminate dams, waterways and drains with bait or its used containers. DO NOT use this product to control native animals. The product is restricted to use in and around buildings (within 2 metres) or within enclosed spaces. In areas

where possums and other non-target animals frequent, baits should be secured inside proper bait stations or bait stations constructed from drainpipes or boxes.

STORAGE AND DISPOSAL

Store in original container, tightly closed, in a safe place. Triple or preferably pressure rinse containers before disposal. Dispose of rinsings and unused or untaken bait in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit. Empty containers and product should not be burnt.

Search for dead mice and rats, burn or bury them, do not place in refuse bins or rubbish tips. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased. Containers that have held bait should not be used for any other purpose.

SAFETY DIRECTIONS

Poisonous if swallowed. Will irritate the skin. May irritate the eyes. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. If on skin and after each baiting, wash thoroughly with soap and water. Use scoop or measure. Vitamin K1 (phytomenadione) is antidotal. When opening the container and using the product, wear elbow-length rubber gloves. After each day's use, wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26. Vitamin K1 (phytomenadione) is antidotal.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

NOTE TO PHYSICIANS AND VETERINARIANS

Difenacoum, the active constituent of Sorex[®] Pro Rodenticide Blocks, is a long lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

This product as supplied is of a high grade and believed to be suitable for any purpose for which it is expressly recommended and must be used in accordance with the Directions for Use given on the label. No responsibility is accepted in respect of the product save those non-excludable conditions implied by any Federal and State legislation or law of a Territory.

® Sorex is a registered trademark of Sorex Limited

® Bitrex is a registered trademark of Macfarlan Smith Limited.

APVMA Approval No. 61842/10KG/0407

Batch No.

D.O.M.

LABL01097



SOREXA[®] Pro
RODENTICIDE BLOCKS



APPROVED LABEL

File name:	61842_41643_10KG_Label_MPL_V01.pdf
Dimensions:	341mm x 179mm
Scale:	70% on A4
Date:	28th May 2007