

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

MOUSEOFF®

ZINC PHOSPHIDE BAIT



Active Constituent: 25g/kg ZINC PHOSPHIDE

For the control of heavy infestations of mice in agricultural situations.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR SCHEDULE 7 POISONS

Net contents: 125kg

DIRECTIONS FOR USE
RESTRAINTS

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).
DO NOT apply bait in a trail.
DO NOT apply bait if heavy rain is imminent.
DO NOT apply to the outer 50m of crop or within 50m of native vegetation. A 50m buffer zone in crop must be employed to reduce risk to birds and other non-target animals.
DO NOT apply bait unless monitoring of mouse numbers indicate the potential for crop damage of economic significance.

Situation	Pest	Rates	Critical Comments
Grain crops, legume crops, canola, safflower, nut crops and pasture. (Note: for all situations -only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seed have been sown into crop stubble or other similar ground cover).	Mice	1kg/ha to achieve an even coverage of 2-3 grains/m^2	Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 grains/m2. It is imperative that all attempts are made to apply bait as evenly as possible to maximise product efficacy and minimise risks to non target animals. Mouseoff Zinc Phosphide Bait will provide a high level of control in most situations. However in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Retreatment after 14 days may therefore be required in certain circumstances. It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods for monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT IS TOO HAZARDOUS TO USE IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIOD

DO NOT harvest any crops or allow livestock to graze baited areas for 14 days after application.

PROTECTION OF WILDLIFE, THE ENVIRONMENT AND LIVESTOCK

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

STORAGE AND DISPOSAL

The contents of each container should be used within 3 months of opening. Store in the closed original container in a dry cool well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition.
Dispose of used containers by crushing and burying below one metre in a disposal pit specifically marked and set up for this purpose, clear of waterways, vegetation and roots. Excess or unused baits must be buried below one metre. Empty containers and product must not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames - forms toxic gas. When opening the container and using baits-wear elbow length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each days use wash gloves and respirator and if rubber, wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone: 13 11 26).
Do not give direct mouth-to-mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy-viva or one way mask. Resuscitate in a well-ventilated area.

MATERIAL SAFETY DATA SHEET:

Additional information is listed in the MSDS which can be obtained from the supplier.

Batch Number:

NRA Approval No. 50532/0800

Date of manufacture:

ANIMAL CONTROL TECHNOLOGIES PTY LTD 46 - 50 Freight Drive, SOMERTON VIC 3062 AUSTRALIA