Dragnet® DUST
Insecticidal Powder
Industrial Strength

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ACTIVE CONSTITUENT:
20 g/kg PERMETHRIN 40:60

NET CONTENTS:
10 kg

GROUP	3A	INSECTICIDE

For the control of various insect pests in accordance with the Directions for Use Table.

SAFETY DIRECTIONS
Will irritate the eyes and skin. Avoid contact with eyes. **DO NOT** inhale dust. When opening the container and using the product wear rubber gloves and a disposable dust mask. After each days use wash gloves and contaminated clothing. Wash hands after use.

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET
Additional information is listed in the Safety Data Sheet, which is available from the supplier or at www.FMCaustralasia.com.au.

GENERAL INSTRUCTIONS

Dragnet® Dust - Insecticidal Powder is ready to use, no further diluent is required. Leave powder undisturbed for as long as possible.

RESISTANCE WARNING
For insecticide resistance management Dragnet® Dust is a Group 3A Insecticide. Some naturally occurring insect biotypes resistant to Dragnet® Dust and other Group 3A Insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Dragnet® Dust or other Group 3A Insecticides are used repeatedly. The effectiveness of Dragnet® Dust on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, FMC accepts no liability for any losses that may result from the failure of Dragnet® Dust to control resistant insects. Dragnet® Dust may be subject to specific resistance management strategies. For further information contact your local supplier, FMC representative or local agricultural department agronomist.

PRECAUTIONS
Avoid contact with food, food utensils, or places where food is prepared or stored. Before use, remove or cover all exposed foodstuffs. Cover all dishes and utensils, and places where food is prepared or stored.

DO NOT apply dust directly on humans.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
Toxic to bees and reptiles. Harmful to fish. **DO NOT** contaminate ponds, waterways or ditches with the chemical or used containers.

PROTECTION OF PETS AND LIVESTOCK
Cover or remove any open food or water containers. Cover or remove fish ponds or aquariums etc. before dusting.

STORAGE AND DISPOSAL
Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple or preferably pressure rinse containers before disposal. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

WARRANTY
FMC makes no warranty expressed or implied, concerning the use of this product other than that indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/ or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

ADDITIONAL GHS STATEMENT(S)
Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves/protective clothing/eye protection/face protection. IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention.

DIRECTIONS FOR USE:

SITUATION		TARGET INSECTS	RATE	CRITICAL COMMENTS
Domestic, Institutional & Industrial premises	Food preparation and storage areas. Living and sleeping areas. Roof, floor, and wall voids and crawl spaces.	Cockroaches, Silverfish, Spiders	5 g/m² applied with a powder blower	Blow powder into cracks and crevices. Pay particular attention to warm, dark corners around cookers, fireplaces, radiators, stoves, pipes, rubbish chutes and places where liquid sprays may not be used. May also be used in difficult to access areas including roof voids, wall cavities, basements, around electrical equipment and behind cupboards.
		Ants		Sprinkle the powder along runs, on and around nests. If Ants are invading domestic premises apply powder to routes by which they enter. Repeat as necessary.
		Fleas & Carpet beetles		Dust powder over floor, brushing into carpet pile and blowing into areas where larvae are found.
		Bedbugs		Apply powder to bedsteads, bedsprings, mattresses, floor coverings, upholstered furniture, cracks in walls, behind torn wall paper, joints in woodwork and other harbourages.
Skin and hide preparation		Hide beetles *(Dermestes ater)*		a) Place skins on drying racks and apply dust using a manual or motorised duster. b) Dust exposed edges of dried, baled skins.
Mushroom beds		Mushroom Sciarid flies		Dust mushroom beds no more frequently than twice a week to control adult and larval flies. Discontinue dusting when mushrooms reach the size of a 10 cent piece.
Seed potatoes		Potato moths *(Phthorimaea operculella)*	300 g/tonne or 15 g/50 kg bag of Seed potatoes	Dust thoroughly immediately after harvest while filling storage containers. Treated seed potatoes are to be used only for sowing; **DO NOT** use for food or feed purposes. Treated seed potatoes should be stored apart from other potatoes and bags or containers should be clearly marked to identify that the contents have been treated.
Large wasp nests		European wasps *(Vespula germanica)*	65 to 130 g per nest	Use hand duster and apply powder through entrance holes (65-130 g/nest according to size of nest and number of entrance holes). Treat at night wherever possible to avoid being stung. Operators should wear clothing protective against the wasp including long-sleeved overalls buttoned at wrist and neck, gloves, hat and bee veil.
Bee nests	In wall cavities	Feral honey bee		
Roof voids of houses which have harboured nesting birds		Bird mites (*Ornithonyssus* spp.)	5 g/m²	Cover roof void areas using a powder blower. Pay particular attention to areas where there is evidence of bird nests. Remove nests and bird-proof roof voids to prevent recurrence.
Trees and stumps in urban areas		Subterranean termites (*Coptotermes* spp.)	125 g per tree	Upon locating nest in tree or stump, drill 3 holes with a maximum diameter of 12 mm. Apply dust directly to the nest using a carbon dioxide gas cylinder and trigger gun. After treatment holes should be coated carefully with a fungicide, e.g. copper naphthenate, and sealed with an appropriate caulking compound.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL, UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD
MUSHROOMS
DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

APVMA Approval No. 52336/10KG/0307

The FMC logo and Dragnet® are Trademarks or Registered trademarks of FMC or its affiliates.

FMC Australasia Pty Ltd.
Level 2, Building B
12 Julius Ave
North Ryde NSW 2113
Phone: 1800 066 355
www.fmcaustralasia.com.au.

SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111
ALL HOURS – AUSTRALIA WIDE