CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS

Racumin®

RAT AND MOUSE PASTE

Active Constituent: 0.37 g/kg COUMATETRALYL

A ready-to-use palatable sachet bait for the control of rats and mice

GENERAL INSTRUCTIONS

Do not apply directly on or near exposed food or food utensils, food preparation areas, livestock feed or water, food storage areas, or onto packaging materials or in areas where such packaging is stored. Do not feed the bait to wild or domestic birds.

Baiting Strategies

Eliminate, as far as practicable, all alternative food sources. Place the bait sachets in areas of known rodent activity and under cover (eg. Behind appliances and furniture, in ceiling voids etc). Inspect baits daily. If eaten quickly increase the number of bait sachets. Continue observation and replenishment until no more bait is taken. Baiting for at least 2 weeks is necessary to reduce mouse or rat numbers. Caution is required in use of this product in and near piggeries. Do NOT allow to contaminate pig food.

PRECAUTION

Should not be used in areas accessible to children.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Place baits in locations inaccessible to other animals and birds. This product is hazardous to pigs, cats and dogs. Do not contaminate ponds, waterways or drains with the product, bait, refuse from the bait stations, or used containers.

STORAGE AND DISPOSAL

Store in the tightly closed, original container in a dry, well-ventilated area, as cool as possible out of direct sunlight. Destroy untaken baits. Any spillage of bait should be removed immediately. Break, crush, or puncture and bury empty containers in a local authority landfill. If not available bury the container below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots.

SAFETY DIRECTIONS

Poisonous if swallowed. If on skin, and after each baiting, wash thoroughly with soap and water. Vitamin K1 (Phytomenadione) is antidotal.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131126.
Additional information is listed in the Material Safety Data Sheet.

LIMITATION OF LIABILITY

This product is designed only for the particular purposes indicated on this label. Failure to use this product strictly as directed may be illegal; prove dangerous; render the product ineffective and avoid any Warranty or Liability of Bayer CropScience Pty Ltd.

Racumin® is a trademark of Bayer
Bitrex™ is a trademark of Macfarlan Smith Limited

NRA Approval No. 51508/0500

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111

DIRECTIONS FOR USE

Restraints

Do NOT place baits in open unless in bait stations.
Do NOT place baits in any position accessible to children, livestock, or domestic pets.
Do NOT remove bait from the paper sachet.

(10 g sachets)

SITUATION	PEST	RATE	CRITICAL COMMENTS
Domestic, industrial, wharf areas, around farm buildings, dairies, meatworks (non-product areas)	Mice and Rats	3-6 sachets per bait site	Place bait sachets in areas of known rodent activity. For rats place a minimum of 3 sachets (30g) per placement. Do not exceed 3m between baits for mice, or 9m for rats. Follow the **Baiting Strategy** detailed below.

(100g sachets)

SITUATION	PEST	RATE	CRITICAL COMMENTS
Domestic, industrial, wharf areas, around farm buildings, dairies, meatworks (non-product areas)	Mice and Rats	1 sachet per bait site	Place bait sachets in areas of known rodent activity. Do not exceed 3 m between baits for mice, or 9 m for rats. Follow the **Baiting Strategy** detailed below.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.