Main panel

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Chipco® GT

FUNGICIDE

ACTIVE CONSTITUENT: 240 g/L IPRODIONE
SOLVENT: 333 g/L LIQUID HYDROCARBONS

GROUP	2	FUNGICIDE

Contents

5 L

For control of fungal diseases in ornamentals and recreational turf as specified in the Directions For Use table.

IMPORTANT: Read the attached booklet before use.

Front page of booklet

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CHIPCO® GT FUNGICIDE

ACTIVE CONSTITUENT: 240 g/L IPRODIONE
SOLVENT: 333 g/L LIQUID HYDROCARBONS

For control of fungal diseases in ornamentals and recreational turf as specified in the Directions For Use table.

STORAGE AND DISPOSAL
Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.
Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.
If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. DO NOT re-use empty containers for any other purpose.

SAFETY DIRECTIONS
Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET
Additional Information is listed in the Safety Data Sheet which is available at www.bayeres.com.au.

Exclusion of Liability
This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty. Ltd. accepts no liability or responsibility for loss or damage from failure to follow such directions and instructions.

IMPORTANT: READ THIS BOOKLET BEFORE USE.

APVMA Approval No.: 50954/58169

Chipco® is a Registered Trademark of the Bayer Group.



Bayer CropScience Pty. Ltd.
A.B.N. 87 000 226 022
391-393 Tooronga Rd
Hawthorn East VIC 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6803
Technical enquiries: 1800 804 479
Website: www.bayeres.com.au

For 24 hour specialist advice in an emergency only phone 1800 033 111

Batch Number: DOM: Barcode

Inside booklet

DIRECTIONS FOR USE

SITUATION	DISEASE	RATE	CRITICAL COMMENTS
Ornamentals	Botrytis blight (*Botrytis cinerea*)	215 ml / 100 L water	Spray at 14 day intervals commencing when the disease first becomes apparent and continue until conditions no longer favour the disease. Spraying Saintpaulia, Poinsettia and Cyclamen flowers may result in some petal scorch. Tepid water should be used and wet plants protected from direct sunlight. This use is subject to an AFRAC anti-resistance strategy: DO NOT apply more than 3 consecutive **Chipco® GT** or related dicarboximide fungicides.
Recreational turf	Brown patch (*Rhizoctonia solani*)	130 to 200 ml / 10 to 150 L water / 100 m^2	Curative program: Repeat treatment after 14 days. Use the higher rate under conditions of severe disease pressure. For preventative treatment: Apply as a monthly spray program. This use is subject to an AFRAC anti-resistance strategy. DO NOT apply more than 2 consecutive sprays of **Chipco® GT** or related dicarboximide fungicides unless tank mixed with a fungicide from a different activity group.
	Dollar spot (*Sclerotinia homoeocarpa*)	130 ml / 10 to 150 L of water / 100 m^2	For curative treatment: Repeat treatment in 14 days. For preventative treatment: Apply monthly. This use is subject to an AFRAC anti-resistance strategy. DO NOT apply more than two consecutive sprays of **Chipco® GT** or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Fusarium patch (*Microdochium nivale*, *Fusarium nivale*) (NOT Qld)	200 ml / 10 to 150 L of water / 100 m^2	
	Spring dead spot (*Leptosphaeria* spp.)	100 ml / 10 to 150 L of water / 100 m^2	For preventative treatment only: Apply as a monthly soil drench throughout the year. Water in thoroughly after application. This use is subject to an AFRAC anti-resistance strategy. DO NOT apply more than two consecutive sprays of **Chipco® GT** or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Helminthosporium disease affecting leaf, crown and roots: (black and white *Helminthosporium* spot, *Bipolaris* spp., *Drechslera* spp., *Exserohilum* spp.)		For curative treatment: Apply as a foliar spray and repeat application after 7 to 14 days. For white spot symptoms on couch grass (*Cynodon dactylon*) only, use higher water volumes of 100 to 150L of water per 100m^2. For preventative treatment: Apply as a high volume foliar spray. Repeat applications at monthly intervals from February to May. This use is subject to an AFRAC anti-resistance strategy. DO NOT apply more than two consecutive sprays of **Chipco® GT** or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Note: Chipco® GT may be very unstable in conditions where the pH is 7 or higher. It is therefore essential to check the pH of the spray solution before adding **Chipco® GT**. A suitable registered buffering agent may have to be added to bring pH down below 7.

Application: Good control requires good coverage. Application should be made using sufficient water to ensure thorough coverage.

Turf nutrition:
This product acts very rapidly and where the turf is of a low nutritional status, a slight yellowing may be seen. This will be corrected by nitrogenous fertiliser application. Where possible, it is recommended that the fertiliser program should precede the fungicide application.

Compatibility:
With any mixture, thoroughly agitate immediately before applying. The use of a surfactant or spray oil is not recommended with **Chipco® GT** as it may result in damage to sensitive plants. DO NOT mix with fertilisers. Mixtures with some fertilisers, e.g. urea, may cause foliar damage. Mixing **Chipco® GT** with **Chipco® Signature Systemic Fungicide** may result in some settling out.
As formulations of other manufacturers' products are beyond the control of Bayer CropScience, all mixtures should be tested prior to mixing commercial quantities.

FUNGICIDE RESISTANCE STRATEGY

GROUP	2	FUNGICIDE

For fungicide resistance management **Chipco® GT** is a Group 2 fungicide. Some naturally occurring individual fungi resistant to **Chipco® GT** and other Group 2 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by **Chipco® GT** and other Group 2 fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience accepts no liability for any losses that may result from the failure of **Chipco® GT** to control resistant fungi.
For further information on managing fungicide resistance contact your local supplier or Bayer Environmental Science representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

DO NOT graze treated areas or feed turf clippings from treated areas to animals including poultry.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.
Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.
If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. DO NOT re-use empty containers for any other purpose.

SAFETY DIRECTIONS
Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET
Additional Information is listed in the Safety Data Sheet which is available at www.bayeres.com.au.

Exclusion of Liability
This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty. Ltd. accepts no liability or responsibility for loss or damage from failure to follow such directions and instructions.

APVMA Approval No.: 50954/58169

Chipco® is a Registered Trademark of the Bayer Group.

Bayer CropScience Pty. Ltd.
A.B.N. 87 000 226 022
391-393 Tooronga Rd
Hawthorn East, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6803
Technical enquiries: 1800 804 479
Website: www.bayeres.com.au

For 24 hour specialist advice in an emergency only phone 1800 033 111